

ICD-10 indications for Fetal Echocardiogram

INDICATION	CODE
FETAL FACTORS	
Arrhythmia - bradycardia (< 120 bpm), persistent	036.83_#
Arrhythmia - heart block, suspected	036.83_#
Arrhythmia - irregular cardiac rhythm, frequent episodes or persistent	036.83_#
Arrhythmia - tachycardia (> 180 bpm), persistent	036.83_#
Cardiac function abnormality, suspected	035.8XX#
Cardiac structural anomaly, suspected	035.8XX#
Chromosomal abnormality (invasive test or NIPT)	035.1XX#
Effusions**	035.8XX#
Hydrops fetalis	036.2_X#
Major extracardiac anomaly (CNS)	035.0XX#
Major extracardiac anomaly (non-CNS)	035.8XX#
Nuchal translucency ≥ 3.5 mm or ≥ 99%ile; (NT 3.0-3.4* or > 95%ile**)	028.3
Placenta or umbilical cord abnormalities (e.g. single umbilical artery)**	043.19_
Systemic venous anomaly* (e.g. persistent right UV, left SVC, absent DV)	035.8XX#
Triplet gestation with ≥ 2 monochorionic	030.11_
Triplet gestation with ≥ 2 monoamniotic	030.12_
Twin gestation - conjoined twins	030.02_
Twin gestation - monochorionic-monoamniotic twins	030.01_
Twin gestation - monochorionic-diamniotic twins	030.03_
MATERNAL or FAMILIAL DISEASE or MATERNAL ENVIRONMENTAL EXPOSURE	
Autoimmune disease with anti-SSA (or SSB**) with (or without*) a prior affected fetus	099.891 & R76.8
Diabetes, gestational, early diagnosis†, A1	024.410
Diabetes, gestational, early diagnosis†, A2, on oral agent	024.415
Diabetes, gestational, early diagnosis†, A2, on insulin	024.414
Diabetes, gestational, early diagnosis†, unspecified control	024.419
Diabetes, pregestational, Type 1	024.01_
Diabetes, pregestational, Type 2	024.11_
First- (or second-*) degree relative with CHD	035.2XX#
First- or second-degree relative with CHD-associated Mendelian inheritance disorder	035.2XX#
In vitro fertilization, including intracytoplasmic sperm injection	009.81_
Phenylketonuria (unknown status or periconceptional phenylalanine level > 10 mg/dL)	099.28_ & E70.0
Rubella infection in the first trimester	035.3XX# & B06 series
Teratogen exposure (e.g. ACE-inhibitor*, carbamazepine*, lithium*, paroxetine*, retinoid)	035.5XX#
Teratogen exposure (e.g. anticonvulsants, NSAID, vitamin A)**	035.5XX#

*: indicates consider

**: AHA-specific guidance

†: diagnosis in the first or early second trimester, (hemoglobin A1c > 6%**).

4th, 5th, or 6th character () is for trimester; use 2 for second trimester and 3 for third trimester. 7th character (#) is for fetal identification; use 0 for singleton or not specified, 1 for fetus 1, 2 for fetus 2, 3 for fetus 3, etc. and 9 for other fetus.